

6046
 6047

NUMBER OF ODD BINOMIAL COEFFICIENTS

HEIKO HARBORTH

ABSTRACT. Let $F(n)$ denote the number of odd numbers in the first n rows of Pascal's triangle, and $\theta = (\log 3)/(\log 2)$. Then $\alpha = \limsup F(n)/n^\theta = 1$, and $\beta = \liminf F(n)/n^\theta = 0.812\ 556\ \dots$

It is known that almost all binomial coefficients are even numbers (see for example [1]-[3]). This means

$$\lim_{n \rightarrow \infty} F(n) / \binom{n+1}{2} = \lim_{n \rightarrow \infty} F(n)/n^2 = 0,$$

if $F(n)$ denotes the number of odd numbers in the first n rows of Pascal's triangle. Recently in [4] and [5] it is asked more precisely for the asymptotic behavior of $F(n)$. Let

$$(1) \quad \alpha = \limsup_{n \rightarrow \infty} F(n)/n^\theta, \quad \beta = \liminf_{n \rightarrow \infty} F(n)/n^\theta,$$

and

$$(2) \quad \theta = (\log 3)/(\log 2) = 1.584\ 962\ \dots$$

Then it is shown in [5] that

$$1 \leq \alpha \leq 1.052, \quad \text{and} \quad 0.72 \leq \beta \leq (9/7)(3/4)^\theta \leq 0.815.$$

Furthermore it is conjectured that 1 and $(9/7)(3/4)^\theta = 3^\theta/7 = 0.814\ 931\ \dots$ are the true values of α and β . In this note we will prove $\alpha = 1$ and $\beta = 0.812\ 556\ \dots$

THEOREM 1. $\alpha = 1$.

PROOF. Since

$$\binom{n}{0} = \binom{n}{n} = 1, \quad \text{and} \quad \binom{n}{i} \equiv 0 \pmod{2}, \quad 1 \leq i \leq n-1,$$

for $n = 2^r$, $r = 0, 1, \dots$, we have the recursion

$$(3) \quad F(2^r + x) = F(2^r) + 2F(x), \quad 0 \leq x \leq 2^r, \quad r = 0, 1, \dots,$$

if, in addition, $F(0) = 0$ is defined. From (3), by induction on r , we get

$$(4) \quad F(2^r) = 3^r,$$

and thus $F(2^r)/2^{r\theta} = 3^r/2^{r\theta} = 1$ for all r , which yields $\alpha \geq 1$.

Next we assert

Received by the editors March 19, 1976.

AMS (MOS) subject classifications (1970). Primary 10L10, 10A30.

Here the reader may recognize the well-known result (see [5] for references) that the number of odd $\binom{n}{r}$ is 2^t , where t is the number of binary digits in n . We insert (9) and (6) in (8), and substitute $2n_r = a$ and $2^t/(3F(n_r)) = b$ to get

$$1 + b \geq \left(1 + \frac{1}{a}\right)^\theta = 1 + \frac{\theta}{a} + \frac{\theta(\theta - 1)}{2a^2} + \theta(\theta - 1) \sum_{i=1}^{\infty} \left(\frac{-1}{a}\right)^{i+2} \frac{(2 - \theta) \cdots (i + 1 - \theta)}{(i + 2)!},$$

$$1 - b \geq \left(1 - \frac{1}{a}\right)^\theta = 1 - \frac{\theta}{a} + \frac{\theta(\theta - 1)}{2a^2} + \theta(\theta - 1) \sum_{i=1}^{\infty} \left(\frac{1}{a}\right)^{i+2} \frac{(2 - \theta) \cdots (i + 1 - \theta)}{(i + 2)!}.$$

Addition of the last two inequalities yields the contradiction

$$2 \geq 2 + \theta(\theta - 1)/a^2 + \cdots > 2.$$

Thus the inequalities (8) cannot both be true, which proves the Lemma.

Now $q_r > 0$ together with the Lemma proves the convergence of $\{q_r\}$. It follows that

$$(10) \quad B \leq q = \lim_{r \rightarrow \infty} q_r < q_{19} = 0.812\ 556 \dots,$$

with

$$n_{19} = 710\ 317 \\ = 2^{19} + 2^{17} + 2^{15} + 2^{14} + 2^{12} + 2^{10} + 2^9 + 2^7 + 2^5 + 2^3 + 2^2 + 1.$$

We still have to prove

$$(11) \quad F(n)/n^\theta > 0.812\ 556 = \gamma.$$

This is true for $1 \leq n \leq 2$, and we assume the validity of (11) for $1 \leq n \leq 2^r$. To obtain the step from r to $r + 1$ in a proof of (11) by induction on r we have to conclude from this assumption that (11) also holds for $n = 2^r + x$, $1 \leq x \leq 2^r$. We divide this interval into eleven intervals:

$$n = 2^{r-s}m + x, \quad 1 \leq x \leq 2^{r-s}, \\ m = n_s \quad \text{for } s = 1, 3, 6, 8, 10, \\ m = n_s - 1 \quad \text{for } s = 2, 4, 5, 7, 9, 10.$$

Let t be the sum of the binary digits of m , and $2^s < m < 2^{s+1}$. Then for $1 \leq x \leq 2^{r-s}$ we get from (3) and (4) that

$$(12) \quad \frac{F(2^{r-s}m + x)}{(2^{r-s}m + x)^\theta} = \frac{3^{r-s}F(m) + 2^tF(x)}{(2^{r-s}m + x)^\theta} > \frac{3^{r-s}F(m) + 2^t\gamma x^\theta}{(2^{r-s}m + x)^\theta} = f_s(x).$$

The unique extremum of $f_s(x)$ is a minimum at

$$x_{\min} = 2^{r-s} (F(m)/\gamma m 2^t)^{1/(\theta-1)}.$$

For $m = n_s$ and $s = 1, 3, 6, 8, 10$ we check by calculation that

$$(13) \quad f_s(x) \geq f_s(x_{\min}) = \left((F(m)/m^\theta)^{1/(1-\theta)} + (\gamma 2^r)^{1/(1-\theta)} \right)^{1-\theta} > \gamma$$

is fulfilled. For $m = n_s - 1$ and $s = 2, 4, 5, 7, 9, 10$ we ascertain that in these cases $x_{\min} > 2^{r-s}$. Then for $s \neq 10$,

$$f_s(x) \geq f_s(2^{r-s}) = \frac{F(n_s - 1) + \gamma 2^{s-1}}{n_s^\theta} = \frac{F(n_s) - (1 - \gamma) 2^{s-1}}{n_s^\theta} > \gamma$$

is seen to be true by calculation. In the case $m = n_{10} - 1$, $s = 10$, we first have

$$f_{10}(x) \geq f_{10}(2^{r-11}) = \frac{3F(n_{10}) - (3 - \gamma) 2^{r-11}}{(2n_{10} - 1)^\theta} > \gamma, \quad 1 \leq x \leq 2^{r-11}.$$

For the remaining partial interval

$$n = 2^{r-10}(n_{10} - 1) + 2^{r-11} + x = 2^{r-11}(2n_{10} - 1) + x, \quad 1 \leq x \leq 2^{r-11},$$

we choose $m = 2n_{10} - 1$ and $s = 11$ in (12), and check the validity of (13).

Now the induction on r is complete, and we have proved (11) for all n . Inequalities (10) and (11) then yield Theorem 2.

At the end we remark that q from (10) probably will be the exact value of β . Moreover, we conjecture for all r ,

$$F(n)/n^\theta \geq q_r \quad \text{for } 2^r \leq n \leq 2^{r+1}.$$

It seems, however, that for a general proof we should know some more properties of the sequence of plus and minus signs beginning with (7). Are there any regularities in this sequence?

REFERENCES

1. N. J. Fine, *Binomial coefficients modulo a prime*, Amer. Math. Monthly **54** (1947), 589-592. MR **9**, 331.
2. H. Harborth, *Über die Teilbarkeit im Pascal-Dreieck*, Math.-Phys. Semesterber. **22** (1975), 13-21.
3. D. Singmaster, *Notes on binomial coefficients. III: Any integer divides almost all binomial coefficients*, J. London Math. Soc. (2) **8** (1974), 555-560.
4. K. B. Stolarsky, *Digital sums and binomial coefficients*, Notices Amer. Math. Soc. **22** (1975), A-669. Abstract #728-A7.
5. ———, *Power and exponential sums of digital sums related to binomial coefficient parity*, SIAM J. Appl. Math. (to appear).

INSTITUT FÜR MATHEMATIK, TECHNISCHE UNIVERSITÄT BRAUNSCHWEIG,
D3300 BRAUNSCHWEIG, WEST GERMANY