

Pattern-Avoiding Permutations

STEVEN FINCH

April 27, 2006

Let $\sigma = \sigma_1\sigma_2\cdots\sigma_m$ be a permutation on $\{1, 2, \dots, m\}$. Define a **pattern** $\tilde{\sigma}$ to be the string $\sigma_1\varepsilon_1\sigma_2\varepsilon_2\cdots\varepsilon_{m-1}\sigma_m$, where each ε_j is either the dash symbol - or the empty string. For example,

1-3-2, 1-32, 132

are three distinct patterns. The first is known as a **classical pattern** (dashes in all $m - 1$ slots); the third is also known as a **consecutive pattern** (no dashes in any slots). Some authors call $\tilde{\sigma}$ a “generalized pattern” and use the word “pattern” exclusively for what we call “classical patterns”.

Let $\tau = \tau_1\tau_2\cdots\tau_n$ be a permutation on $\{1, 2, \dots, n\}$, where $n \geq m$. We say that τ **contains** $\tilde{\sigma}$ if there exist $1 \leq i_1 < i_2 < \dots < i_m \leq n$ such that

- for each $1 \leq j \leq m - 1$, if ε_j is empty, then $i_{j+1} = i_j + 1$;
- for all $1 \leq k \leq m$, $1 \leq l \leq m$, we have $\tau_{i_k} < \tau_{i_l}$ if and only if $\sigma_k < \sigma_l$.

The string $\tau_{i_1}\tau_{i_2}\cdots\tau_{i_m}$ is called an **occurrence** of $\tilde{\sigma}$ in τ . If τ does not contain $\tilde{\sigma}$, then we say τ **avoids** $\tilde{\sigma}$ or that τ is **$\tilde{\sigma}$ -avoiding**. For example,

24531 contains 1-3-2

because 253 has the same relative order as 132, but

42351 avoids 1-3-2.

As another example,

6725341 contains 4132

because 7253 has the same relative order as 4132 and consists of four consecutive elements, but

41352 avoids 4132.

As a final example,

3542716 contains 12-4-3

⁰Copyright © 2006 by Steven R. Finch. All rights reserved.

because 3576 has the same relative order as 1243 and its first two elements are consecutive, but

$$3542716 \text{ avoids } 12\text{-}43.$$

Define $\alpha_n(\tilde{\sigma})$ to be the number of n -symbol, $\tilde{\sigma}$ -avoiding permutations. We naturally wish to understand the rate of growth of $\alpha_n(\tilde{\sigma})$ with increasing n .

0.1. Classical Patterns. The Stanley-Wilf conjecture, proved by Marcus & Tardos [1], was rephrased by Arratia [2] as follows:

$$L(\tilde{\sigma}) = \lim_{n \rightarrow \infty} (\alpha_n(\sigma_1\text{-}\sigma_2\text{-}\dots\text{-}\sigma_m))^{1/n}$$

exists and is finite. We have [3, 4, 5, 6, 7]

$$L(\tilde{\sigma}) = 4 \quad \text{when } m = 3,$$

$$L(1\text{-}2\text{-}\dots\text{-}m) = (m - 1)^2 \quad \text{for all } m \geq 2,$$

$$L(1\text{-}3\text{-}4\text{-}2) = 8,$$

$$L(1\text{-}2\text{-}4\text{-}5\text{-}3) = (1 + \sqrt{8})^2 = 9 + 4\sqrt{2}.$$

A conjecture that $L(\tilde{\sigma}) \leq (m - 1)^2$ has been disproved [8]:

$$9.35 \leq L(1\text{-}3\text{-}2\text{-}4) \leq 288$$

and hence the maximum limiting value (as a function of m) remains open. Also, we wonder if $L(\tilde{\sigma})$ is always necessarily an algebraic number.

0.2. Consecutive Patterns. Elizalde & Noy [9, 10] examined the cases $m = 3$ and $m = 4$. The quantities $\alpha_n(123)$ and $\alpha_n(132)$ satisfy

$$\alpha_n(123) \sim \gamma_1 \cdot \rho_1^n \cdot n!, \quad \alpha_n(132) \sim \gamma_2 \cdot \rho_2^n \cdot n!$$

where

$$\rho_1 = 3\sqrt{3}/(2\pi) = 0.8269933431\dots, \quad \gamma_1 = \exp\left(\pi/(3\sqrt{3})\right) = 1.8305194665\dots,$$

$$\rho_2 = 1/\xi = 0.7839769312\dots, \quad \gamma_2 = \exp(\xi^2/2) = 2.2558142944\dots$$

and $\xi = 1.2755477364\dots$ is the unique positive solution of

$$\int_0^x \exp(-t^2/2) dt = 1, \quad \text{that is,} \quad \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{x}{\sqrt{2}}\right) = 1.$$

The quantities $\alpha_n(1342)$, $\alpha_n(1234)$ and $\alpha_n(1243)$ satisfy

$$\alpha_n(1342) \sim \gamma_1 \cdot \rho_1^n \cdot n!, \quad \alpha_n(1234) \sim \gamma_2 \cdot \rho_2^n \cdot n!, \quad \alpha_n(1243) \sim \gamma_3 \cdot \rho_3^n \cdot n!$$

where

$$\begin{aligned} \rho_1 &= 1/\xi = 0.9546118344\dots, & \gamma_1 &= 1.8305194\dots, \\ \rho_2 &= 1/\eta = 0.9630055289\dots, & \gamma_2 &= 2.2558142\dots, \\ \rho_3 &= 1/\zeta = 0.9528914198\dots, & \gamma_3 &= 1.6043282\dots; \end{aligned}$$

ξ , η and ζ are the smallest positive solutions of

$$\int_0^x \exp(-t^3/6) dt = 1, \quad \cos(y) - \sin(y) + \exp(-y) = 0,$$

$$3^{1/2} \int_0^z \text{Ai}(-s) ds + \int_0^z \text{Bi}(-s) ds = \frac{3^{1/3}\Gamma(1/3)}{\pi},$$

respectively, where $\text{Ai}(t)$ and $\text{Bi}(t)$ are the Airy functions [11].

0.3. Other Results. Elizalde [12, 13] proved that

$$\lim_{n \rightarrow \infty} \left(\frac{\alpha_n(1-23-4)}{n!} \right)^{1/n} = 0$$

and believed that the same applies to $\alpha_n(12-34)$, although a proof is not yet known. Ehrenborg, Kitaev & Perry [14] gave more detailed asymptotic expansions for $\alpha_n(123)$ and $\alpha_n(132)$; a similar “translation” of combinatorics into operator eigenvalue analysis was explored in [15]. The field is wide open for research.

Let us focus on classical patterns in the following. Define $\sigma \leq \tau$ if τ contains $\tilde{\sigma}$. A **permutation class** C is a set of permutations such that, if $\tau \in C$ and $\sigma \leq \tau$, then $\sigma \in C$. Let C_n denote the permutations in C of length n . If $C = \{\text{all permutations}\}$, then $|C_n| = n!$; such behavior is regarded as degenerate and this case is excluded from now on. The Marcus-Tardos theorem implies that, for nondegenerate C ,

$$L(C) = \limsup_{n \rightarrow \infty} |C_n|^{1/n} < \infty.$$

Consider the set R of all growth rates $L(C)$ and the derived set R' of all accumulation points of R . Vatter [16] proved that

$$\inf \{r \in R : r > 2\} = 2.0659948920\dots$$

which is the unique positive zero of $1 + 2x + x^2 + x^3 - x^4$, and

$$\inf \{s : s \text{ is an accumulation point of } R'\} = 2.2055694304\dots$$

which is the unique positive zero of $1 + 2x^2 - x^3$. Albert & Linton [17] proved that R is uncountable and thus contains transcendental numbers. Vatter [18] subsequently proved that

$$\inf \{t : R \text{ contains the interval } (t, \infty)\} \leq 2.4818728574\dots$$

which is the unique positive zero of $-1 - 2x - 2x^2 - 2x^4 + x^5$ and conjectured that \leq can be replaced by $=$. The question of whether \limsup in the definition of $L(C)$ can be replaced by \lim is also unanswered.

REFERENCES

- [1] A. Marcus and G. Tardos, Excluded permutation matrices and the Stanley-Wilf conjecture, *J. Combin. Theory Ser. A* 107 (2004) 153–160; MR2063960 (2005b:05009).
- [2] R. Arratia, On the Stanley-Wilf conjecture for the number of permutations avoiding a given pattern, *Elec. J. Combin.* 6 (1999) N1; MR1710623 (2000e:05005).
- [3] A. Regev, Asymptotic values for degrees associated with strips of Young diagrams, *Adv. Math.* 41 (1981) 115–136; MR0625890 (82h:20015).
- [4] M. Bóna, Exact enumeration of 1342-avoiding permutations: a close link with labeled trees and planar maps, *J. Combin. Theory Ser. A* 80 (1997) 257–272; math.CO/9702223; MR1485138 (98j:05003).
- [5] M. Bóna, The limit of a Stanley-Wilf sequence is not always rational, and layered patterns beat monotone patterns, *J. Combin. Theory Ser. A* 110 (2005) 223–235; math.CO/0403502; MR2142176 (2005k:05005).
- [6] M. Bóna, New records on Stanley-Wilf limits, submitted (2006); available online at <http://www.math.ufl.edu/~bona/cikkek.html>.
- [7] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A0022558, A005802, A061552.
- [8] M. H. Albert, M. Elder, A. Rechnitzer, P. Westcott and M. Zabrocki, On the Wilf-Stanley limit of 4231-avoiding permutations and a conjecture of Arratia, math.CO/0502504.

- [9] S. Elizalde and M. Noy, Consecutive patterns in permutations, *Adv. Appl. Math.* 30 (2003) 110–125; available online at <http://www.math.dartmouth.edu/~sergi/>; MR1979785 (2004d:05004).
- [10] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A049774, A111004, A113228, A113229, A117156, A117158, A117226.
- [11] S. R. Finch, Airy function zeroes, unpublished note (2004).
- [12] S. Elizalde, Asymptotic enumeration of permutations avoiding generalized patterns, *Adv. Appl. Math.* 36 (2006) 138–155; math.CO/0505254.
- [13] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A113226, A113227.
- [14] R. Ehrenborg, S. Kitaev and P. Perry, A spectral approach to pattern-avoiding permutations, *Formal Power Series and Algebraic Combinatorics (FPSAC)*, Proc. 2006 San Diego conf., http://garsia.math.yorku.ca/fpsac06/en/contrib_papers.html
- [15] R. Ehrenborg and Y. Farjoun, Asymptotics of the Euler number of bipartite graphs, arXiv:0704.1782.
- [16] V. Vatter, Small permutation classes, arXiv:0712.4006.
- [17] M. H. Albert and S. A. Linton, Growing at a perfect speed, <http://www.circa.mcs.st-and.ac.uk/pre-prints.html#2007>.
- [18] V. Vatter, Permutation classes of every growth rate (a.k.a. Stanley-Wilf limit) above 2.48187..., arXiv:0807.2815.