

Integers sequences A328348 and A328350 to A328356

Pierre-Alain Sallard

October 14, 2019

Abstract

The sequence A328348 is the building block for the calculation of the sums of positive integers whose decimal notation only uses two distinct, non-zero digits.

The sequences A328350 to A328356 are also building blocks for the calculation of the sums of positive integers, in the case where more than two distinct, non-zero digits are used .

1 Sums of whole numbers whose decimal notation only uses two distinct, non-zero digits

Let α and β be two distinct non-zero digits (i.e. $(\alpha, \beta) \in \{1, 2, \dots, 9\}^2$, with $\alpha \neq \beta$), let $n \in \mathbb{N}$, and let $E_{\alpha, \beta, n}$ be the set of whole numbers whose decimal notation only uses the α and β digits and that have at most n such digits. Formally,

$$E_{\alpha, \beta, n} = \left\{ x \in \mathbb{N} \mid \exists p \in \llbracket 0; n-1 \rrbracket, \exists (a_0, \dots, a_p) \in \{\alpha, \beta\}^{p+1}, x = \sum_{i=0}^p a_i 10^i \right\}$$

We define

$$S_{\alpha, \beta, n} = \sum_{x \in E_{\alpha, \beta, n}} x$$

We state that

$$S_{\alpha, \beta, n} = (\alpha + \beta) \times u_{2, n}$$

where

$$\forall n \in \mathbb{N}, u_{2, n} = \frac{10 \cdot 20^n - 19 \cdot 2^n + 9}{171}$$

The sequence $(u_{2, n})_{n \in \mathbb{N}} = (0, 1, 23, 467, 9355, 187131, 3742683, 74853787, 1497075995, \dots)$ corresponds to the A328348 OEIS sequence.

For example, with $\{\alpha, \beta\} = \{1, 2\}$ and $n = 3$, we have

$$E_{1, 2, 3} = \{1, 2, 11, 12, 21, 22, 111, 112, 121, 122, 211, 212, 221, 222\}$$

and

$$S_{1,2,3} = \sum_{x \in E_{1,2,3}} x = (1+2) \times u_3 = 1401.$$

Proof : by definition, $S_{\alpha,\beta,n} = \sum_{p=0}^{n-1} \sum_{(a_0,\dots,a_p) \in \{\alpha,\beta\}^{p+1}} \sum_{i=0}^p a_i 10^i$. For a given $p \in$

$\llbracket 0; n-1 \rrbracket$, a given $\epsilon \in \{\alpha, \beta\}$ and a given $j \in \llbracket 0; p \rrbracket$, $\text{Card}(\{(a_0, \dots, a_{j-1}, \epsilon, a_{j+1}, \dots, a_p) \mid (a_i)_{i \neq j} \in \{\alpha, \beta\}^p\}) = 2^p$: it means that the digit α appears 2^p times in first position ($j = 0$), 2^p times in second position ($j = 1$), etc. Therefore,

$$\forall p \in \llbracket 0; n-1 \rrbracket, \sum_{(a_0,\dots,a_p) \in \{\alpha,\beta\}^{p+1}} \sum_{i=0}^p a_i 10^i = \alpha \times 2^p \times \sum_{i=0}^p 10^i + \beta \times 2^p \times \sum_{i=0}^p 10^i =$$

$$(\alpha + \beta) 2^p \sum_{i=0}^p 10^i = (\alpha + \beta) \frac{10 \cdot 20^p - 2^p}{9}. \text{ Thus } S_{\alpha,\beta,n} = \frac{\alpha + \beta}{9} \sum_{p=0}^{n-1} (10 \cdot 20^p - 2^p) =$$

$$\frac{\alpha + \beta}{9 \times 19} (10 \cdot 20^n - 19 \cdot 2^n + 9). \quad \square$$

Remark : it can be noted that, in the expression of $S_{\alpha,\beta,n}$, only the sum $\alpha + \beta$ matters and not the individual values of α and β . As a consequence, as soon as $\alpha + \beta = \alpha' + \beta'$, we have $S_{\alpha,\beta,n} = S_{\alpha',\beta',n}$. For example, for all positive integer n , $S_{1,7,n} = S_{2,6,n} = S_{3,5,n}$.

2 Generalisation : sums of whole numbers whose decimal notation only uses k distinct, non-zero digits

Instead of selecting only two distinct digits, lets take $k \in \llbracket 2; 9 \rrbracket$ distinct, non-zero digits and consider the set $E_{\alpha_1, \dots, \alpha_k, n}$ of whole numbers whose decimal notation only uses $\alpha_1, \dots, \alpha_k$ as digits and that have at most n such digits.

We define

$$S_{\alpha_1, \dots, \alpha_k, n} = \sum_{x \in E_{\alpha_1, \dots, \alpha_k, n}} x$$

With a proof similar to the above, it can be established that

$$S_{\alpha_1, \dots, \alpha_k, n} = \left(\sum_{i=1}^k \alpha_i \right) \times u_{k,n}$$

where

$$\forall n \in \mathbb{N}, u_{k,n} = \frac{10(k-1)(10k)^n - (10k-1)k^n + 9}{9(k-1)(10k-1)}$$

The sequences appearing in the above formula are :

- for $k = 3$, $(u_{3,n})_{n \in \mathbb{N}} = (0, 1, 34, 1033, 31030, 931021, 27930994, 837930913, \dots)$ corresponds to the A328350 OEIS sequence.
- for $k = 4$, $(u_{4,n})_{n \in \mathbb{N}} = (0, 1, 45, 1821, 72925, 2917341, 116695005, 4667805661, \dots)$ corresponds to the A328351 OEIS sequence.

- for $k = 5$, $(u_{5,n})_{n \in \mathbb{N}} = (0, 1, 56, 2831, 141706, 7086081, 354307956, 17715417331, \dots)$ corresponds to the A328352 OEIS sequence.
- for $k = 6$, $(u_{6,n})_{n \in \mathbb{N}} = (0, 1, 67, 4063, 244039, 14643895, 878643031, 52718637847, \dots)$ corresponds to the A328353 OEIS sequence.
- for $k = 7$, $(u_{7,n})_{n \in \mathbb{N}} = (0, 1, 78, 5517, 386590, 27064101, 1894506678, 132615604717, \dots)$ corresponds to the A328354 OEIS sequence.
- for $k = 8$, $(u_{8,n})_{n \in \mathbb{N}} = (0, 1, 89, 7193, 576025, 46086681, 3686971929, 294958053913, \dots)$ corresponds to the A328355 OEIS sequence.
- for $k = 9$, $(u_{9,n})_{n \in \mathbb{N}} = (0, 1, 100, 9091, 819010, 73718281, 6634711720, 597124652671, \dots)$ corresponds to the A328356 OEIS sequence.

For example, with $k = 5$, $\{\alpha_1, \dots, \alpha_5\} = \{1, 2, 3, 4, 5\}$ and $n = 2$, we have

$$E_{1,2,3,4,5,2} = \{1, 2, 3, 4, 5, 11, 12, 13, 14, 15, 21, 22, 23, 24, 25, \dots, 54, 55\}$$

and

$$S_{1,2,3,4,5,2} = (1 + 2 + 3 + 4 + 5) \times u_{5,2} = 15 \times 56 = 840$$